

Detection of Change in Supercycles in ER Ursae Majoris

Polina ZEMKO,^{1,2} Taichi KATO,² Sergei Yu. SHUGAROV,^{1,3}

¹ *Sternberg Astronomical Institute, Moscow University, Universitetsky Ave., 13, Moscow 119992, Russia*
polina.zemko@gmail.com

² *Department of Astronomy, Kyoto University, Kyoto 606-8502*
tkato@kustastro.kyoto-u.ac.jp

³ *Astronomical Institute of the Slovak Academy of Sciences, 05960, Tatranska Lomnica, the Slovak Republic*

(Received 200 0; accepted 200 0)

Abstract

We examined data on about 20 years of observations of ER UMa available in AAVSO, VSNET, AFOEV, NSVS, VSOLJ databases together with published light curves. The obtained $O - C$ diagram revealed a systematic change of the supercycle (time interval between two successive superoutbursts) within 43.6 and 59.2 d. The time-scale of this change variation is from 300 to ~ 1900 d. The number of normal outbursts within the supercycles also varied from 4 to 6 although no strong correlation between this number and supercycle length was found. We suggest that appearance of negative superhumps is responsible for the observed variations of number of normal outbursts. Our results generally confirms the expectations by the thermal-tidal instability theory.

Key words: accretion, accretion disks — stars: dwarf novae — stars: individual (ER UMa) — stars: novae, cataclysmic variables

1. Introduction

Dwarf novae (DNe) are a class of cataclysmic variables (CVs) — close binary systems consisting of a white dwarf as a primary component and a red or brown-dwarf secondary transferring matter via the Roche-lobe overflow. Because of the instabilities that take place in the accretion disks normal outbursts and less frequent superoutbursts with larger amplitudes take place (see Warner 1995, Hellier 2001). DNe that show both normal outbursts and superoutbursts are called the SU UMa-type stars. These stars have one distinguishing property — light-curve modulation with the periods close to, but different from, the orbital one that appear near the superoutburst maxima and sometimes persist at other times (Vogt 1980). These modulations are called positive superhumps if their period is longer than the orbital one and negative if shorter. Positive superhumps are believed to arise from the periodic viscous dissipation in a disk driven to resonant oscillation by the tidal field of the secondary star, and precessing slowly in the prograde direction (see Whitehurst 1988, Hirose, Osaki 1990, Lubow 1991, Smith et al. 2007, Wood et al. 2011) and negative superhumps arise from the precession of tilted accretion disk, respectively (Wood et al. 2009, Montgomery, Martin 2010).

SU UMa stars have two extreme subgroups with very long (about decades) and short (tens of days) supercycles. They are WZ Sge and ER UMa subtypes, respectively. There are five ER UMa-type stars known by now: RZ LMi (Nogami et al. 1995), DI UMa (Kato et al. 1996), ER UMa (Kato, Kunjaya 1995), V1159 Ori (Robertson et al. 1995), IX Dra (Ishiooka et al. 2001) and all of them show extremely short supercycle (18.9 – 54 d) and posi-

tive superhumps (Olech et al. 2008). Moreover, ER UMa-type stars were known to show a pronounced stability of the supercycle length and outburst pattern, which can be particularly seen in the folded light-curves and $O - C$ diagrams presented in (Robertson et al. 1995). This stability and short supercycles can be explained as a result of a very high and constant mass-transfer rate in the framework of thermal tidal instability (TTI) model (Osaki 1995a).

However, long-term monitoring of V1159 Ori revealed secular changes in the supercycle and a hint of the periodic behavior of such variations (Kato 2001). The TTI model suggests that this phenomenon is the consequence of a change in mass-transfer rate in the binary system, although the origin of these changes is still an open question. The hibernation scenario, or the theory of cyclic evolution, suggested by Livio (1992), also assumes that mass-transfer rate in CVs can vary secularly and proposes irradiation of the secondary by the white dwarf which is still hot after the nova eruption as one of the possible mechanisms of an increased mass-transfer. However, the time-scale of $\dot{M}$ variation in this scenario is very long comparing to these observations. On the other hand for V1159 Ori, Kato (2001) proposed that solar-type cycles of the secondary which are possibly observed in cataclysmic variables (Bianchini 1988, Ak et al. 2001) may be responsible for the phenomenon.

Here we report that another one ER UMa-type star — ER UMa itself, also shows variations of the supercycle. A discussion on the origin of this phenomenon is even more intriguing in the context of recent discovery by Ohshima et al. (2012) of persistent negative superhumps in this dwarf nova. The presence of negative superhumps was not taken into account in past studies of SU UMa-type

stars, while this factor may strongly affect the outburst behavior. Osaki, Kato (2012) have also found that the presence of negative superhumps is correlated with normal outbursts, thus it is reasonable to investigate the possibility whether the presence of negative superhumps is related to the long-term variations of the outburst behavior.

A comprehensive study of long-term variations in ER UMa-type stars is therefore essential for better understanding the evolution of dwarf novae in general, in particular changes in mass transfer-rate, the mechanisms that lead to the appearance of negative superhumps and their effect on the outburst behavior.

2. Observation and Data Analysis

2.1. Light-curves

We analyzed all the data on ER UMa available in American Association of Variable Stars Observers (AAVSO)¹, Variable Stars NETwork (VSNET)², Association Française des Observateurs d'Etoiles Variables (AFOEV)³, Northern Sky Variability Survey (NSVS)⁴ and Variable Star Observers League in Japan (VSOLJ)⁵ databases which represent more than 20 years of observations. The earliest superoutburst was on 1992 April 27 (JD2448740) and the latest one on 2012 June 12 (JD2456090.7). We also used observations obtained by one of the authors (S. Shugarov) in the Stará Lesná Observatory of Slovak Academy of Sciences, and by the group of E. Pavlenko in the Crimean Astrophysical Observatory. A long-term light-curve of ER UMa from these observations is presented in figure 1.

2.2. Moments of the superoutbursts and $O-C$ diagram

In order to follow the supercycle change in ER UMa we plotted $O-C$ diagram using the data mentioned. The moments of the superoutbursts and normal outbursts were determined visually from the light curves. The mean period essential for plotting such diagram was determined by summing up the cycle counts and was chosen to be 48.7 d by the least-squares method. There were several gaps in observations usually 3-4 cycles long, particularly around the solar conjunction, which can be also seen in figure 1. The time intervals between two subsequent detected superoutbursts in some such cases were large enough to bring considerable uncertainties in calculations of the number of cycles in such gaps. For every uncertain moment of the superoutburst we checked the two closest possible values of the cycle number, for each of them calculated the mean period for the few closest supercycles before, across and after the gap and regarded the variant as less probable if it led to the significant period jump. The list of the moments of superoutbursts, corresponding supercycle count — E (the number of superoutbursts passed since JD2448740), supercycle length,

measured by the linear fitting of successive five epochs of $O-C$ diagram and the number of normal outbursts observed within the supercycle is presented in the table 1.

2.3. Normal outbursts phases

We noticed that ER UMa shows two types of outburst behavior with 3-4 normal outbursts within the supercycle observed and with 5 or more. Examples of these types can be seen in figure 2 and figure 3. Following Osaki, Kato (2012) and Smak (1985) here we use Type S for the stages with many normal outbursts and short intervals and Type L with reduced numbers of normal outbursts and long intervals. Because of the gaps in observations and because the data were not very precise for the most part of supercycles it was possible to determine only approximate numbers of normal outbursts within them, or even only lower limits. In the table 1 we give the probable numbers, as “3-4” for the cases where it was impossible to choose the unique value, and lower limits, as “>4”, for the supercycles, where for example four normal outbursts were detected but the significant part of the supercycle remained unobserved, and we could not say whether there were other normal outbursts or not.

3. Discussion

3.1. $O-C$ Behavior, Comparison with Other SU UMa DNe

The $O-C$ diagram for the whole analyzed period is presented in figure 4. We can see that ER UMa underwent several stages with a relatively stable supercycle length (linear portion in the diagram) which lasted for 6 – 30 cycles and terminated by an abrupt period change. The local mean supercycle length was determined by the linear fit of each time interval with relatively stable supercycle in the $O-C$ diagram. It was shown that the supercycle varied from 43.7 at the very beginning of the $O-C$ diagram to 59.2 d. The local values of the supercycle length are also presented in the table 1.

The $O-C$ behavior presented in figure 4 is reminiscent to that of V1159 Ori, shown in figure 2. in (Kato 2001). Although one can see such a pattern, for example between $E=60$ and $E=125$ cycles in figure 4, the time-scale of the variability is significantly longer in ER UMa. A detailed analysis of the supercycle variations in SU UMa-type stars was presented by (Vogt 1980). Although only ordinary SU UMa-stars were discussed in Vogt (1980) (we do not consider a comparison with SS Cyg-type DNe since it is out of the scope of this paper), the published $O-C$ diagrams have the same features as ER UMa, particularly of SU UMa itself (figure 1 in Vogt 1980). Supercycles of SU UMa remained stable for relatively long periods of time (10 – 20 cycles) and switched randomly between several characteristic values. Nevertheless, while Vogt (1980) has shown that the supercycles in SU UMa possess 3 characteristic values, in ER UMa there are no preferable supercycle lengths as can be seen in figure 5.

¹ <http://www.aavso.org/>

² <http://www.kusastro.kyoto-u.ac.jp/vsnet/>

³ <http://cdsarc.u-strasbg.fr/afoev/>

⁴ <http://skydot.lanl.gov/nsvs/nsvs.php>

⁵ <http://vsolj.cetus-net.org/>




Fig. 1. Long-term light-curve of ER UMa from AAVSO, VSNET, AFOEV, NSVS and VSOLJ observations.




Fig. 4. $O - C$ diagram for ER UMa with the phases of normal outbursts and the types of superhumps observed. Open and filled circles correspond to L and S normal outburst phases, respectively, supercycles with undetermined phases are marked with crosses.




Fig. 5. Distribution of the supercycle lengths of ER UMa.




Fig. 2. Example of the L type normal outburst behavior of ER UMa (March - June 2011)

Fig. 3. Example of the S type normal outburst behavior of ER UMa (January - May 1995)

3.2. Supercycle Length Change

We measured the local lengths of the supercycle by the linear fitting of $O - C$ diagram and plotted them against time. Results can be seen in figure 6 where the supercycle length change is presented. Supercycle lengths (T_s) appeared to change discontinuously and show oscillations superimposed on a secular increasing trend. The rate of supercycle growth is $dT_s/dE = 0.033(6)$ d/cycle and the period derivative is $dT_s/T_s = 6.7(6) \times 10^{-4}$, respectively. We used a half-delete jackknife method described in (Shao, Tu 1995) to determine reliability of the obtained trend. It yielded that the period increase is statistically significant with a 90% confidence. Nevertheless, these formal error estimates should not be taken seriously. In Otulakowska-Hypka et al. (2012) it was shown that the period derivative for IX Dra is 1.8×10^{-3} , comparable with that found for ER UMa. We suppose that it is a typical value of supercycle change rate for ER UMa-type stars. In figure 5 there is a hint of bimodal distribution of the periods of supercycle with maxima at 48 and 53 d, respectively. Osaki (1995a) proposed a model of ER UMa type stars based on the TTI theory which successfully explained the light-curves and short recurrence times of the superoutbursts assuming extremely high mass-transfer rate (about 10 times higher than that expected from CV evolutionary scenario based on gravitational-wave radiation). It also provided a dependence between the period of supercycle and mass-transfer rate (presented in figure 1 in Osaki 1995a) and the minimum possible supercycle length for ER UMa-type stars (about 40 d). According to this dependence there are two possible values of $\dot{M}$ for each T_s . These two possibilities can be distinguished by considering the duty cycle of superoutbursts (ratio of the duration of superoutburst to the supercycle length). In our case, the possibility of larger $\dot{M}$ can be excluded because the observed duty cycles were much smaller than 0.5. Observed variations in the supercycle length of ER UMa correspond to the change of mass-transfer rate from $3.8 \dot{M}_{16}$ to $2 \dot{M}_{16}$ where $\dot{M}_{16}$ is the mass-transfer rate in units of 10^{16} g s^{-1} and the increasing trend in T_s means that the mass-transfer rate in this system is decreasing secularly. It is also remarkable that the most significant change of the supercycle and the mass-transfer rate (almost twice) occurred in first break in the $O - C$ diagram (JD2449932) right after the initial long stable interval. All the further supercycle changes were less dramatic. The period varied between 46.2 and 54.1 d which corresponds to a $\sim 15\%$ change in mass-transfer rate. It is also notable that the minimum value of the supercycle was not below the shortest achievable supercycle in (Osaki 1995a) and we therefore do not need to assume an additional mechanism as in RZ LMi (Osaki 1995b)

3.3. Normal Outbursts

Another important problem in the behavior of ER UMa is the correlation between the length of the supercycle and the number of normal outbursts within it. For the systems with a high mass-transfer rate (like ER UMa), both




Fig. 6. Variation of the supercycle length measured by the linear fitting of the $O - C$ diagram. The solid line shows the secular increasing trend, whose statistical significance was verified by a half-delete jackknife method (Shao, Tu 1995) with a 90% confidence. Open and filled circles correspond to L and S normal outburst phases, respectively, supercycles with undetermined phases marked with crosses.

the waiting time for normal outbursts (T_n) and the waiting time for superoutbursts (T_s) are approximately proportional to $\dot{M}^{-1}$ (Osaki 1995b). In this case T_n/T_s does not depend on $\dot{M}$ and the number of normal outbursts between two successive superoutbursts should be constant according to the model of ER UMa stars proposed by Osaki (1995a). However, as was mentioned in Subsection 2.3, ER UMa shows two types of normal outburst phases: a phase with 3-4 normal outbursts observed and with more than 4. In a Type L phase presented in the figure 2 ER UMa underwent four normal outbursts (marked with arrows) that occurred once in 7 days. In figure 3 five normal outbursts can be seen, moreover in the second supercycle the 6-th normal outburst near JD2449824 was likely unobserved because of the gap in the observations. The time intervals between two subsequent normal outbursts were 4 days. In figure 4 and figure 6, filled circles correspond to the supercycles with Type S normal outburst phase and open circles to the supercycles with Type L phase. If it was impossible to determine the type of the phase within the supercycle we mark them with crosses. It can be seen from these plots that there are no strong correlation between the normal outburst phases and the length of the supercycle. Types L and S were observed in any value of the supercycle length. Bends on the $O - C$ diagram, or shifts of the supercycle period, did not affect the number of normal outbursts. Apparently there should be some additional factor, which is responsible for the change of the

normal outbursts phases. Osaki, Kato (2012), Ohshima et al. (2012) suggested that the type of phase may depend on the absence or presence of negative superhumps. This possibility is discussed in the following subsection.

3.4. Negative Superhumps and Outburst Behavior

Negative superhumps are believed to arise from the precession of a tilted accretion disk, nevertheless the origin of the disk tilt is still under debate. In such cases the hot spot appears not only at the disks rim but deeper in the primaries potential, too. Montgomery, Martin (2010) proposed a mechanism based on the hydrodynamic lift. Differences between the supersonic velocities of the accretion flow under and below the disk cause the appearance of the lift force and make the disk tilted when the mass transfer is high. The accretion stream can thus reach the inner parts of the accretion disk. Interaction of the magnetic fields with rotating plasma was also suggested as a possible source of the disk tilt (Murray, Armitage 1998, Murray et al. 2002, Smak 2009). A disk tilt is likely to affect the outburst behavior of SU UMa stars, and it is very important to study the correlation between the presence of negative superhumps (as an indicator of the tilt of the disks) and outbursting activity. In 2011 it became evident that persistent negative superhumps in ER UMa were observed even during the superoutburst (Ohshima et al. 2012) while a retrospective search for the literature and the available data suggests that they were ab-

sent in the past. Kato, Kunjaya (1995) in their discovery paper reported the detection of positive superhumps in ER UMa. No negative superhumps which are considered as characteristic feature of ER UMa in the present state, were observed by them in 1994. The first possible detection of negative superhumps in ER UMa was claimed by Gao et al. (1999), but the amplitudes were smaller than in Ohshima et al. (2012) and it is likely that, even if the negative superhumps did exist in 1998, they were not so strong in 1998 as in 2011. Since then the presence of the superhumps with period less than the orbital one was confirmed by Zhao et al. (2006), Kjurkchieva, Marchev (2010). In our retrospective study we also confirmed the presence of negative superhumps in 2010 in the AAVSO data. It appears likely that the negative superhumps were present since 2006.

3.5. Correlation with Normal Outburst Phase

TTI theory suggests that a mass-transfer to a tilted disk is expected to reduce the number of normal outbursts (Osaki, Kato 2012). Since the accretion stream in such cases can reach the inner parts of the disk, material is accumulating not primarily on the disks rim. It leads to the reduction of “outside-in” outbursts. Here we examine the correlation between the normal outburst phase and the presence of negative superhumps. In the figure 4 we mark the types of superhumps observed by the authors mentioned above. One can see that the presence of negative superhumps seems to depend strongly on the phase of the normal outbursts. The first possible detection of negative superhumps was near the $E = 44$ on figure 4 when the first L stage was observed. The next one, claimed by Zhao et al. (2006) was on 2004 February 29 which corresponds to the $E = 88$, just at the transition from S to L phases. Negative superhumps observed by Kjurkchieva, Marchev (2010) in 2010 were in $E = 119$ at the diagram, also at the beginning of L phase. According to the detailed light curves of ER UMa provided by the observational campaign of the VSNET collaboration (Kato et al. 2004) negative superhumps have been observed since the 2011 ($E = 135$) and four normal outbursts occurred within each supercycle. It confirms that the phase of the normal outbursts can be strongly affected by the presence of negative superhumps, and negative superhumps can indeed reduce the number of normal outbursts. So we can now assume that Type L phase can be regarded as “negative superhump” phase.

3.6. Correlation with the Supercycle length

According to figure 4, positive and negative superhumps were observed both on the descending and ascending branches of the $O - C$ diagram, and they do not correlate with the supercycle length. Negative superhumps were with no doubt observed since 2011. During this period $O - C$ underwent a bend near $E = 150$ but it had no affect on the presence of negative superhumps. In the TTI model, the next superoutburst occurs when the disk stores a certain amount of angular momentum (J_{crit}), and supercycle is not expected to change if $\dot{J}_{crit}$ from the sec-

ondary is constant regardless of the point of accretion on the disk. A disk tilt therefore is not expected to affect the supercycle length and our analysis confirmed this expectation.

4. Conclusions

In this paper we examined data of about 20 years of observations of ER UMa. We followed the evolution of the change in supercycle length, the behavior of normal outbursts and the types of superhumps observed. We found the oscillations of the supercycle length superimposed with a secular increasing trend and associated it with a gradual change of the mass-transfer rate in the binary system. The estimated period derivative was $dT_s/T_s = 6.7(6) \times 10^{-4}$. We also found that ER UMa shows two types of normal outburst phases with 3-4 and with 5 or more normal outbursts within the supercycle. We also examined the possible correlation between the supercycle length, normal outburst phase and the presence of negative superhumps and found a strong dependence between the last two. A reduced number of normal outbursts was observed when negative superhumps were detected. It confirms the assumption of Osaki, Kato (2012) that negative superhumps tend to suppress the normal outbursts. No correlation between the length of the supercycle and other characteristics was found.

This work was supported by the Bilateral International Exchange Program (2012) offered by the Kyoto University and Global COE “The Next Generation of Physics, Spun from Universality and Emergence” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

Dr. Shugarov acknowledge the Slovak Academy of Sciences grant VEGA No. 2/0038/10, grant RFBR No. 09-02-00225 and NSH-2374.2012.2 (Russia).

We acknowledge with thanks the variable star observers from the American Association of Variable Stars Observers, Variable Stars NETwork, Association Française des Observateurs d’Etoiles Variables, Northern Sky Variability Survey and Variable Star Observers League in Japan. We also acknowledge the group of Elena Pavlenko for the data presented and would like to thank Yoji Osaki for valuable comments.

References

- Ak, T., Ozkan, M. T., & Mattei, J. A. 2001, A&A, 369, 882
- Bianchini, A. 1988, IBVS, 3136, 1
- Gao, W., Li, Z., Wu, X., Zhang, Z., & Li, Y. 1999, ApJL, 527, L55
- Hellier, C. 2001, Cataclysmic Variable Stars: how and why they vary (Berlin: Springer-Verlag)
- Hirose, M., & Osaki, Y. 1990, PASJ, 42, 135
- Ishioke, R., Kato, T., Uemura, M., Iwamatsu, H., Matsumoto, K., Martin, B. E., Billings, G. W., & Novak, R. 2001, PASJ, 53, L51
- Kato, T. 2001, PASJ, 53, L17
- Kato, T., & Kunjaya, C. 1995, PASJ, 47, 163

- Kato, T., Nogami, D., & Baba, H. 1996, PASJ, 48, L93
- Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2004, PASJ, 56, S1
- Kjurkchieva, D., & Marchev, D. 2010, Publ. de l’Observatoire Astronomique de Beograd, 90, 147
- Livio, M. 1992, in ASP Conf. Ser. 29, Cataclysmic Variable Stars, ed. N. Vogt (San Francisco: ASP), p. 269
- Lubow, S. H. 1991, ApJ, 381, 259
- Montgomery, M. M., & Martin, E. L. 2010, ApJ, 722, 989
- Murray, J. R., & Armitage, P. J. 1998, MNRAS, 300, 561
- Murray, J. R., Chakrabarty, D., Wynn, G. A., & Kramer, L. 2002, MNRAS, 335, 247
- Nogami, D., Kato, T., Masuda, S., Hirata, R., Matsumoto, K., Tanabe, K., & Yokoo, T. 1995, PASJ, 47, 897
- Ohshima, T., et al. 2012, PASJ, 64, L3
- Olech, A., Wiśniewski, M., Złoczewski, K., Cook, L. M., Mularczyk, K., & Kedzierski, P. 2008, Acta Astron., 58, 131
- Osaki, Y. 1995a, PASJ, 47, L11
- Osaki, Y. 1995b, PASJ, 47, L25
- Osaki, Y., & Kato, T. 2012, PASJ, in prss (arXiv astro-ph/1212.1516)
- Otulakowska-Hypka, M., Olech, A., de Miguel, E., Rutkowski, A., Koff, R., & Bąkowska, K. 2012, A&A, in prss (arXiv astro-ph/1211.1924)
- Robertson, J. W., Honeycutt, R. K., & Turner, G. W. 1995, PASP, 107, 443
- Shao, J., & Tu, D. 1995, The Jackknife and Bootstrap (Berlin: Springer-Verlag)
- Smak, J. 1985, Acta Astron., 35, 357
- Smak, J. 2009, Acta Astron., 59, 419
- Smith, A. J., Haswell, C. A., Murray, J. R., Truss, M. R., & Foulkes, S. B. 2007, MNRAS, 378, 785
- Vogt, N. 1980, A&A, 88, 66
- Warner, B. 1995, Cataclysmic Variable Stars (Cambridge: Cambridge University Press)
- Whitehurst, R. 1988, MNRAS, 232, 35
- Wood, M. A., Still, M. D., Howell, S. B., Cannizzo, J. K., & Smale, A. P. 2011, ApJ, 741, 105
- Wood, M. A., Thomas, D. M., & Simpson, J. C. 2009, MNRAS, pp 2110–2121
- Zhao, Y., Li, Z., Wu, X., Peng, Q., Zhang, Z., & Li, Z. 2006, PASJ, 58, 367

Table 1. Moments of the superoutbursts, supercycle lengths (T_s) and the number of normal outbursts within the supercycle (N_{NO}). Since the moments of superoutbursts were determined from several databases, multiple entries for the same E sometimes occur.

E	Date	JD	T_s (d)	N_{NO}	E	Date	JD	T_s (d)	N_{NO}
0	1992/04/27	2448740	49.6		90	2003/12/18	2452992	45.4	3-4
3	1992/09/05	2448871	49.0		91	2004/01/28	2453033	44.0	4
6	1993/01/13	2449001	44.1		91	2004/01/28	2453033	43.3	
7	1993/02/27	2449046	44.5		92	2004/03/13	2453078	46.5	4
8	1993/04/17	2449095	45.0	>4	92	2004/03/13	2453078	46.8	
9	1993/05/29	2449137	43.6	>4	93	2004/05/01	2453127	47.5	4
10	1993/07/12	2449181	44.4		94	2004/06/16	2453173	46.3	3-4
11	1993/08/22	2449222	43.9		94	2004/06/16	2453173	46.6	
12	1993/10/14	2449275	44.0		96	2004/09/15	2453264	47.2	
13	1993/11/18	2449310	43.6		97	2004/11/05	2453315	47.8	
14	1994/01/04	2449357	42.1		97	2004/11/06	2453315	49.2	
15	1994/02/15	2449399	43.1		98	2004/12/25	2453365	48.3	3-4
16	1994/03/29	2449441	42.6	>4	99	2005/02/10	2453412	49.3	3-4
17	1994/05/11	2449484	42.5	>4	99	2005/02/10	2453412	49.8	
18	1994/06/24	2449528	42.4	5-6	100	2005/04/04	2453465	52.1	4
19	1994/08/03	2449568	42.1		101	2005/05/22	2453513	52.2	3-4
20	1994/09/15	2449611	42.1		103	2005/09/08	2453621	52.4	
21	1994/10/27	2449653	43.5		104	2005/10/29	2453673	53.5	
22	1994/12/09	2449696	45.0		104	2005/10/30	2453673	53.4	
23	1995/01/25	2449743	46.9	>3	105	2005/12/23	2453728	54.9	
24	1995/03/14	2449791	47.3	5	105	2005/12/23	2453728	55.2	
25	1995/05/02	2449840	48.71	5	105	2005/12/24	2453728	56.0	
27	1995/08/02	2449932	50.5		106	2006/02/17	2453784	53.3	
28	1995/09/29	2449990	54.4	5	106	2006/02/17	2453784	54.2	4
29	1995/11/24	2450046	59.6		107	2006/04/08	2453834	54.7	6
30	1996/01/31	2450114	59.4		108	2006/06/06	2453893	55.5	
31	1996/03/25	2450168	55.8	>4	108	2006/06/07	2453894	56.1	>5
32	1996/05/22	2450226	48.9		111	2006/11/20	2454060	55.2	
33	1996/07/04	2450269	47.2		111	2006/11/21	2454060	55.0	
36	1996/11/22	2450410	47.0		112	2007/01/13	2454114	54.5	
37	1997/01/06	2450455	47.6	4	112	2007/01/14	2454114	55.0	
38	1997/03/02	2450510	48.1		113	2007/03/10	2454169	54.9	
39	1997/04/15	2450554	47.6	4	113	2007/03/11	2454171	53.7	6
40	1997/06/01	2450601	46.2	4	114	2007/05/03	2454224	53.6	6
42	1997/09/04	2450696	46.0		118	2007/12/03	2454438	52.9	
44	1997/12/03	2450786	45.8	3-4	118	2007/12/04	2454438	51.4	
45	1998/01/16	2450830	44.8	3-4	119	2008/01/20	2454486	45.6	
46	1998/03/05	2450878	44.9	3-4	120	2008/03/03	2454529	46.0	3-4
47	1998/04/15	2450919	46.0		120	2008/03/04	2454530	47.0	
48	1998/06/01	2450966	49.7	5	121	2008/04/21	2454578	49.3	
49	1998/07/21	2451016	52.2		121	2008/04/23	2454580	49.2	3-4
50	1998/09/21	2451078	52.8		122	2008/06/10	2454628	49.7	
51	1998/11/06	2451124	50.6		122	2008/06/11	2454629	49.6	
52	1998/12/28	2451176	49.2		125	2008/11/07	2454778	50.1	3-4
53	1999/02/10	2451220	49.8		125	2008/11/08	2454778	49.8	
54	1999/04/07	2451276	51.6	>4	126	2008/12/29	2454830	49.6	
55	1999/05/24	2451323	52.6	5	127	2009/02/14	2454877	48.9	
57	1999/09/11	2451433	51.0		127	2009/02/15	2454878	48.2	4
58	1999/10/31	2451483	51.0		128	2009/04/04	2454926	51.2	
59	1999/12/13	2451526	47.6	4-5	128	2009/04/05	2454926	47.4	4
60	2000/02/06	2451581	46.0		129	2009/05/29	2454981	47.0	5
61	2000/03/18	2451622	46.2	5	134	2010/01/14	2455211	46.7	
62	2000/04/30	2451665	44.7	5	135	2010/03/01	2455257	45.8	
63	2000/06/19	2451715	46.7	>5	135	2010/03/01	2455257	46.6	
64	2000/08/01	2451758	47.0	>5	136	2010/04/14	2455301	47.7	4

Table 1. (Continued.)

E	Date	JD	T_s (d)	N_{NO}	E	Date	JD	T_s (d)	N_{NO}
65	2000/09/21	2451809	46.5	>5	136	2010/04/20	2455307	46.0	3-4
66	2000/11/04	2451853	47.9		137	2010/06/05	2455353	45.6	
67	2000/12/21	2451900	47.3		141	2010/12/03	2455534	45.2	
68	2001/02/11	2451952	48.4		142	2011/01/16	2455578	45.2	
69	2001/03/27	2451996	48.4	6	143	2011/03/04	2455624	45.8	4
70	2001/05/17	2452047	48.6		143	2011/03/04	2455625	48.1	4
74	2001/11/26	2452240	48.6	6	143	2011/03/05	2455626	49.5	4
75	2002/01/17	2452292	48.0		144	2011/04/22	2455674	49.5	4
76	2002/03/01	2452335	45.0	>4	144	2011/04/23	2455675	49.0	4
77	2002/04/18	2452383	45.1		144	2011/04/24	2455676	48.9	4
81	2002/10/10	2452558	46.1		145	2011/06/11	2455724	48.3	
82	2002/12/02	2452611	46.6		145	2011/06/11	2455724	48.9	
83	2003/01/23	2452663	50.0	>5	147	2011/09/16	2455820	50.6	
84	2003/03/10	2452709	49.2		148	2011/11/07	2455872	51.3	
85	2003/04/29	2452759	48.3		149	2012/01/02	2455929	53.0	
85	2003/04/30	2452760	48.2		149	2012/01/02	2455929	53.9	
86	2003/06/16	2452807	47.9	6-7	150	2012/02/20	2455978	53.4	
88	2003/09/20	2452903	45.9		150	2012/02/27	2455984	53.8	
88	2003/09/21	2452903	45.1		152	2012/06/11	2456089	54.4	
89	2003/10/27	2452940	43.5		152	2012/06/12	2456090	52.8	